


Carbetocin

MADE IN GERMANY

Solution for Injection

100 µg
1 ml



Approved for room temperature storage, providing greater convenience and enhanced flexibility in handling and transport.

Carbetocin EVER Pharma is indicated for the prevention of postpartum haemorrhage due to uterine atony following delivery of the infant, whether by caesarean section or vaginal birth.

- **Room temperature stable** – no requirement for refrigerated storage or cold-chain transport, reducing logistical complexity and lowering storage and transportation costs.
- Available in both **vial and ampoule** presentations to support diverse clinical preferences and practice environments.



www.everpharma.com

 **EVER**
PHARMA

Carbetocin

MADE IN GERMANY

Solution for Injection

Indications	Carbetocin EVER Pharma is indicated for the prevention of postpartum haemorrhage due to uterine atony.
Active Ingredient	Carbetocin
Excipients	Sodium dihydrogenphosphate dihydrate, Sodium acetate trihydrate, Mannitol, Sodium hydroxide (for pH adjustment), Hydrochloric acid (for pH adjustment), Water for injections
Presentations	<ul style="list-style-type: none">■ 100 µg/1 ml solution for injection
Strengths	Each 1 ml of solution contains 100 microgramms of carbetocin in solution.
Stability	<ul style="list-style-type: none">■ Unopened: 24 months. Do not store above 30°C.■ After 1st Opening: Use immediately
Primary Packaging	<ul style="list-style-type: none">■ Type I colourless 1 ml glass ampoule■ Type I colourless glass vial sealed with a halobutyl rubber stopper and an aluminium crimp cap covered with a brown flip off cap.
Pack sizes	1 or 5 ampoules per pack. 1 or 5 vials per pack. Vials may or may not be sheathed in a protective sleeve.



Plastic sleeving for safer handling

Carbetocin EVER Pharma 100 micrograms/ml solution for injection. Composition: 1 ml of solution contains 100 micrograms of carbetocin. List of excipients: Sodium dihydrogenphosphate dihydrate, Sodium acetate trihydrate, Mannitol, Sodium hydroxide (for pH adjustment), Hydrochloric acid (for pH adjustment), Water for injections. Therapeutic indications: For the prevention of postpartum haemorrhage due to uterine atony. Contraindications: During pregnancy and labour before delivery of the infant. Carbetocin must not be used for the induction of labour. Hypersensitivity to carbetocin, oxytocin or to any of the excipients listed in section 6.1. Hepatic or renal disease. Serious cardiovascular disorders. Epilepsy. Side effects: Intravenous administration: Very common: Headache, Tremor, Hypotension, Flushing, Nausea, Abdominal pain, Pruritus, Feeling of warmth. Common: Anaemia, Dizziness, Chest pain, Dyspnoea, Metallic taste, Vomiting, Back pain, Chills, Pain. Not known: Hypersensitivity (including anaphylactic reaction), Tachycardia, Bradycardia which can lead to cardiac arrest, Arrhythmia, Myocardial ischaemia, and QT prolongation. In the clinical trials sweating was reported as sporadic cases. Intramuscular administration: Uncommon: Anaemia, Headache, Dizziness, Tachycardia, Hypotension, Chest pain, Nausea, Abdominal pain, Vomiting, Back pain, Muscular weakness, Chills, Pyrexia, Pain. Rare: Tremor, Flushing, Dyspnoea, Pruritus, Urinary retention. Not known: Hypersensitivity (including anaphylactic reaction), Bradycardia which can lead to cardiac arrest, Arrhythmia, Myocardial ischaemia, and QT prolongation. More information available in the summary of product characteristics. Only available on prescription. Last update: April 2025. Marketing Authorisation Holder: EVER Valinject GmbH, Oberburgau 3, 4866 Unterach am Attersee, Austria.

The product is currently under regulatory review.

CAR/INT/01/2026/1

